



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The grass on 1 acre in 1 week with what grows $= 1 + \frac{1}{9} = \frac{10}{9}$.

$\frac{2}{9} - \frac{1}{9} = \frac{1}{9}$, the amount of the standing grass eaten in 1 week.

$1 \div \frac{1}{9} = 9$ weeks, the time required.

SOME INTERESTING RULES IN MULTIPLICATION.

BY MARY M. CURRIER, WENTWORTH, N. H.

RULE 1. To multiply one number by another, the multiplier consisting of two digits of which the left-hand digit is 1:

Multiply each figure of the multiplicand by the right-hand figure of the multiplier, and to each product add the figure of the multiplicand following the one multiplied.

Example 1. Multiply 1675 by 13.

$$\begin{array}{r} 1675 \times 13 = 21775 \\ 5 \times 3 = 15 \\ 7 \times 3 = 21, \quad 21 + 1 + 5 = 27 \\ 6 \times 3 = 18, \quad 18 + 2 + 7 = 27 \\ 1 \times 3 = 3, \quad 3 + 2 + 6 = 11 \\ \quad \quad \quad 1 + 1 = 2 \end{array}$$

Taking the digits in the units place in these several products beginning with the last, we have, $21775 = 13 \times 1675$.

Example 2. Multiply 40928 by 17.

$$\begin{array}{r} 8 \times 7 = 56 \\ 2 \times 7 = 14, \quad 14 + 5 + 8 = 27 \\ 9 \times 7 = 63, \quad 63 + 2 + 2 = 67 \\ 0 \times 7 = 0, \quad 0 + 6 + 9 = 15 \\ 4 \times 7 = 28, \quad 28 + 1 + 0 = 29 \\ \quad \quad \quad 2 + 4 = 6 \end{array}$$

$\therefore 695776 = 17 \times 40928$.

RULE 2. To multiply one number by another, the multiplier consisting of three digits, of which the two at the left are ones:

Multiply each figure of the multiplicand by the right hand figure of the multiplier and to each product add the two figures following the one multiplied and the digit in the tens place of the preceding product.

Example 1. Multiply 340726 by 114.

$$\begin{array}{r} 6 \times 4 = 24 \\ 2 \times 4 = 8, \quad 8 + 2 + 6 = 16 \\ 7 \times 4 = 28, \quad 28 + 1 + 2 + 6 = 37 \\ 0 \times 4 = 0, \quad 0 + 3 + 7 + 2 = 12 \\ 4 \times 4 = 16, \quad 16 + 1 + 0 + 7 = 24 \\ 3 \times 4 = 12, \quad 12 + 2 + 4 + 0 = 18 \\ \quad \quad \quad 1 + 3 + 4 = 8 \\ \quad \quad \quad 3 = 3 \end{array}$$

$\therefore 38842764 = 114 \times 340726$.

RULE 3. To multiply one number by another, the multiplier consisting of two digits of which the left hand digit is 2.

Multiply each figure of the multiplicand by the right-hand figure of the multiplier and to each product add twice the figures of the multiplicand following the one multiplied and the digit in tens place of the preceding product.

Example 1. Multiply 3495013 by 26.

$$\begin{array}{rcl}
 3 \times 6 & & = 18 \\
 1 \times 6 = 6, & 6 + 1 + 2 \times 3 = & 13 \\
 0 \times 6 = 0, & 0 + 1 + 2 \times 1 = & 3 \\
 5 \times 6 = 30, & 30 + 0 + 2 \times 0 = & 30 \\
 9 \times 6 = 54, & 54 + 3 + 2 \times 5 = & 67 \\
 4 \times 6 = 24, & 24 + 6 + 2 \times 9 = & 48 \\
 3 \times 6 = 18, & 18 + 4 + 2 \times 30 = & 30 \\
 & 3 + 2 \times 3 = & 9
 \end{array}$$

$$\therefore 90870338 = 26 \times 3495013.$$

From these three rules one may see how the principle may be carried further.

NOTE. We deem these rules of sufficient interest to merit publication in this department. It may be seen that they are results of a compact arrangement of the following method: Multiply 3492 by 117.

$$\begin{array}{rcl}
 3492 & 2 \times 7 & = 14 \\
 117 & 9 \times 7 = 63, & 63 + 1 + 2 = 66 \\
 \text{---} & 4 \times 7 = 28, & 28 + 6 + 9 + 2 = 45 \\
 14 & 3 \times 7 = 21, & 21 + 4 + 4 + 9 = 38 \\
 63. & & 3 + 3 + 4 = 10 \\
 28.. & & 1 + 3 = 4 \\
 21... & & \\
 3492. & \therefore 408564 = 117 \times 3492. & \\
 3492.. & & \\
 \text{---} & & \\
 408564 & &
 \end{array}$$

EDITOR.

ALGEBRA.

139. Proposed by JOHN M. COLAW, A. M., Monterey, Va.

Solve neatly $\sqrt[4]{m-x} = \sqrt[4]{n} - \sqrt[4]{x}$.

Solution by J. H. DRUMMOND, LL. D., Portland, Me.

$$(m-x)^{\frac{1}{4}} = n^{\frac{1}{4}} - x^{\frac{1}{4}} \dots (1).$$

Put $n = p^4$ and $x = y^4$. Substituting these values in (1), raising to the fourth power, and reducing, we have

$$y^4 - 2py^3 + 3p^2y^2 - 2py^3 = \frac{m - p^4}{2}.$$

By inspection we see that adding p^4 to the first member it becomes a perfect square. Adding p^4 to both members and extracting the square root we have

$$y^2 - py + p^2 = \pm \sqrt{\frac{m + p^4}{2}}; \text{ then } y = \frac{p \pm [-3p^2 \pm 2\sqrt{(2m + 2p^4)}]^{\frac{1}{2}}}{2}.$$

Restoring values of y and p , we have